

Available online at www.sciencedirect.com

SciVerse ScienceDirect

www.nrjournal.com

Intermittent fasting during Ramadan attenuates proinflammatory cytokines and immune cells in healthy subjects

“Mo’ez Al-Islam” E. Faris^{a,b,*}, Safia Kacimi^c, Ref’at A. Al-Kurd^b,
 Mohammad A. Fararjeh^d, Yasser K. Bustanji^c, Mohammad K. Mohammad^c,
 Mohammad L. Salem^e

^a Department of Clinical Nutrition, College of Applied Medical Sciences, University of Hail, Hail, Kingdom of Saudi Arabia

^b Department of Nutrition, Faculty of Pharmacy and Medical Sciences, Petra University, Amman, Jordan

^c Department of Clinical Pharmacy and Biopharmaceutics, Faculty of Pharmacy, The University of Jordan, Amman, Jordan

^d Faculty of Medical Laboratory Technology, Hashemite University, Zarka, Jordan

^e Zoology Department, Faculty of Science, Tanta University, Tanta, Egypt

ARTICLE INFO

Article history:

Received 21 October 2011

Revised 15 June 2012

Accepted 29 June 2012

Keywords:

Fasting

Caloric restriction

Inflammation

Interleukin-1 β

Interleukin-6

Leukocytes

Tumor necrosis factor α

ABSTRACT

Intermittent fasting and caloric restriction have been shown to extend life expectancy and reduce inflammation and cancer promotion in animal models. It was hypothesized that intermittent prolonged fasting practiced during the month of Ramadan (RIF) could positively affect the inflammatory state. To investigate this hypothesis, a cross-sectional study was designed to investigate the impact of RIF on selected inflammatory cytokines and immune biomarkers in healthy subjects. Fifty (21 men and 29 women) healthy volunteers who practiced Ramadan fasting were recruited for the investigation of circulating proinflammatory cytokines (interleukin [IL]-1 β , IL-6, and tumor necrosis factor α), immune cells (total leukocytes, monocytes, granulocytes, and lymphocytes), and anthropometric and dietary assessments. The investigations were conducted 1 week before Ramadan fasting, at the end of the third week of Ramadan, and 1 month after the cessation of Ramadan month. The proinflammatory cytokines IL-1 β , IL-6, and tumor necrosis factor α ; systolic and diastolic blood pressures; body weight; and body fat percentage were significantly lower ($P < .05$) during Ramadan as compared with before Ramadan or after the cessation of Ramadan fasting. Immune cells significantly decreased during Ramadan but still remained within the reference ranges. These results indicate that RIF attenuates inflammatory status of the body by suppressing proinflammatory cytokine expression and decreasing body fat and circulating levels of leukocytes.

© 2012 Elsevier Inc. All rights reserved.

Abbreviations: BMI, body mass index; CR, caloric restriction; DBP, diastolic blood pressure; HC, hip circumference; HDL, high-density lipoprotein; IF, intermittent fasting; IL-1 β , interleukin-1 β ; IL-6, interleukin-6; LPL, lipoprotein lipase; RIF, Ramadan intermittent fasting; SBP, systolic blood pressure; TG, triglycerides; TNF- α , tumor necrosis factor α ; WBC, white blood cells; WC, waist circumference.

* Corresponding author. Tel.: +966 5 49847886; fax: +966 6 5316982.

E-mail address: moezf@uop.edu.jo (“M.’A.-I.”E. Faris).

0271-5317/\$ – see front matter © 2012 Elsevier Inc. All rights reserved.

<http://dx.doi.org/10.1016/j.nutres.2012.06.021>

1. Introduction

Excessive caloric intake and subsequent obesity are characterized by a chronic state of inflammation including high circulating proinflammatory cytokine levels, a condition that is described as “low-grade inflammation” [1,2]. In this condition, typically 2- to 3-fold increase in the systemic concentrations of tumor necrosis factor- α (TNF- α), interleukin (IL)-1 β , and IL-6 is observed [3]. This increase in the expression of the proinflammatory cytokines may contribute to the induction of autoimmune diseases such as rheumatoid arthritis [4] and inflammatory diseases including atherosclerosis [5], insulin resistance [6], cardiovascular diseases [7], and tissue damage associated with many types of cancer [8].

Over the past few years, an increasing number of physiological effects of intermittent fasting (IF) have been documented in studies on rodents, monkeys, and humans [9]. Prominent among these are increased lifespan [10], decreased mortality from cancers and cardiovascular diseases [11,12], improved insulin sensitivity [13], and reduced oxidative stress and inflammation [14]. Interestingly, the inflammatory biomarkers such as IL-6 and C-reactive protein were found to be significantly depressed by short and long IF [15,16]. In overweight adults, caloric restriction (CR) has improved the clinical findings for patients with moderate asthma and reduced markers of oxidative stress and inflammation [17] and reduced metabolic disease risk markers [18]. Furthermore, it is noteworthy to point out that previous studies have shown that IF reduces oxidative stress and inflammation in tissues, including the brain [19]. Arumugam et al [19] showed that multiple neuroprotective pathways were activated and inflammatory pathways were suppressed by IF in young mice and that aging impaired the ability of IF to modulate these pathways adaptively. In rats, IF and CR have improved cardiovascular and hormonal responses to stress and substantially affect blood pressure and heart rates [20,21].

One of the 5 pillars of Islam is that healthy adult Muslims must fast from dawn to sunset during the holy month of Ramadan. Over the past 40 years, emerging evidence from epidemiologic studies has supported the health-related benefits of Ramadan IF (RIF), including improved insulin sensitivity [22], decreased atherogenic risk, oxidative stress, and inflammation [16,23]. Furthermore, RIF decreased the levels of some circulating cytokines [16,24]. However, there is no available evidence regarding the effect of RIF on circulating IL-1 β , IL-6, and TNF- α and the impact of dietary changes in the people who fast during RIF on proinflammatory cytokines. Therefore, we hypothesized that, compared with nonfasting periods, healthy individuals would show lower levels in the circulating proinflammatory cytokines and immune cells during RIF. To test the hypothesis, we evaluated the anti-inflammatory effect of prolonged (8-24 hours) IF practiced during the holy month of Ramadan in randomly selected healthy volunteers and by using a follow-up model for a cross-sectional study design. The specific objectives of the study were to (i) examine the effect of RIF on the serum levels of the proinflammatory cytokines, namely, IL-1 β , IL-6, and TNF- α ; (ii) elaborate the effects of RIF on leukocyte differential count, total white blood cells (WBC), anthropometric characteristics,

blood pressure, and dietary intake of calorie and antioxidant and energy-yielding nutrients; (iii) determine the correlation between the serum cytokine levels and dietary intakes of calorie, body weight and body fat percentage, systolic (SBP) and diastolic (DBP) blood pressures. Results of the present study are expected to open an avenue for further studies to explore the underlying mechanisms of the anti-inflammatory effect of RIF.

2. Methods and materials

This study was conducted in August to September 2009 over 3 stages: 7 days before Ramadan (T1), after the end of the third week of Ramadan fasting (after 21 days of RIF) (T2), and 1 month after the end of Ramadan fasting month (after 60 days from the start of Ramadan or 30 days from the cessation of Ramadan month) (T3). The range of fasting hours during the month was about 14 to 15 hours a day.

2.1. Subjects

To maintain sample homogeneity and avoid disparities and different confounding cultural and socioeconomic factors, all participants were chosen from the same living community of medium economic level (Rusaifa city of Zarka governorate, the Hashemite Kingdom of Jordan). A total number of 151 volunteers who visited the Al-Quds Medical Laboratories were recruited. The inclusion criteria necessitated that subjects were fasting at least 21 days of the Ramadan month and were not taking any drug or receiving any medical treatment immediately before or during the study. Those who were taking chronic medications, especially those affecting the inflammatory status such as nonsteroidal anti-inflammatory drugs were excluded before beginning of the study. Of the 151 volunteers, only 50 healthy subjects (21 men aged 18-49 years and 29 women aged 18-51 years) were deprived of the exclusion criteria and completed the 3 stages of the study. All enrolled participants were required to provide a written informed consent before starting the study. The study protocol was approved by Petra University Committee on Human Research Ethics. A questionnaire was administered, in which medical history and socioeconomic information were obtained through a personal interview with each subject by a well-trained staff member. It is noteworthy to mention that female subjects did not fast during their menstrual period because Islamic rules do not permit fasting during the menstrual period. Accordingly, the fasting period in female subjects ranged from 25 to 23 days, whereas for men, it was 30 days.

2.2. Blood sampling

Venous blood samples were collected from the enrolled volunteers during the 3 stages or time points. Each subject served as his own control by comparing his/her values before Ramadan (T1) with those during (T2) and after Ramadan (T3). Blood sampling was conducted in the private medical laboratory once a time at T1, T2, and T3 in the same hours

for the 3 time points of the study. It was performed early (first 1-2 hours, 10-12 AM) during the period of investigation (between 10 AM and 4 PM) in each time point. To narrow the time interval for blood sampling and thus to eliminate the effect of day time on blood variables, volunteers were asked to come early (at 10 AM), and blood was first drawn from most of the volunteers according to their arrivals to the laboratory, and then the less time-sensitive assessments (anthropometric and dietary) were subsequently performed during the rest of the investigation period. The obtained sera were separated within an hour by centrifuging the blood samples at room temperature for 5 minutes at 4000 rpm (Sigma 3-18K lab Centrifuge, Osterode am Harz, Germany). Aliquots of the blood samples were directly analyzed for complete blood count cells using Auto Hematology Analyzer (Mindray BC-3000, Shenzhen, China). At each visit, SBP and DBP (mm Hg) were measured using mercury sphygmomanometer, with the subject in a seated position after a 5-minute rest.

2.3. Anthropometric measurements

During each of the 3 visits, anthropometric measurements such as body weight and height (Seca, Hamburg, Germany), waist (WC) and hip (HC) circumferences (nonstretchable measuring tape), and body fat percentage (TANITA body composition analyzer, Tokyo, Japan) were performed by a well-trained staff member. Body weight was measured at the same time to the nearest 0.1 kg; WC and HC, to the nearest 0.01 m; and body height, to nearest 0.01 m. Body mass index (BMI; kg/m²) and waist-to-hip ratio were the calculated. Volunteers with BMI less than 25 kg/m² were considered having normal weight, and those with a BMI of 25 kg/m² or greater were considered having overweight [25].

2.4. Assessment of dietary intakes

There were no special nutritional regimens or dietary recommendations given to the subjects during any stage of the study. Instead, a 24-hour food recall was taken from each subject during the 3 stages. Nutrient intakes were calculated using computerized tables from the Food Processor SQL food and diet analysis software system (version 8.4; Esha Research, Salem, OR, USA).

2.5. Determination of cytokine levels

For cytokine measurements, serum samples were aliquoted, coded, and stored at -80°C until being analyzed. Owing to the insufficient volume of the serum collected, TNF- α level was measured in 48 subjects, whereas IL-1 β and IL-6 levels were measured in only 40 subjects.

Samples from the same volunteer were measured in the same assay to reduce the effect of interassay variation on serum cytokine levels. All samples were freeze-thawed once a time. At the end of the study, the 3 frozen samples drawn during the 3 time points (T1, T2, and T3) for each subject were thawed at the same time and then used in the biochemical measurement by using the same kit and personnel and under the same conditions, to eliminate sources of errors, to avoid the negative impact of cycled freezing-thawing on blood proteins and other components, and to avoid cytokine degradation. In addition, different kits were used within the

same week. Serum levels of TNF- α , IL-1 β , and IL-6 were measured by a sandwich enzyme-linked immunosorbent assay using commercially available matched monoclonal and biotinylated, antibody pairs commercial kits (PeproTech for TNF- α and IL-6, and R&D Systems DuoSet for IL-1 β) (Minneapolis, MN, USA). The measurements were undertaken according to the manufacturer's instructions (DuoSet ELISA Development System; R&D Systems). The measurements of cytokines were performed at the specified absorbance and were detected using a microplate reader (BioTek, Winooski, VT, USA). All measurements were performed in triplicates, and concentrations of these mediators were determined using constructed standard curves.

2.6. Statistical analyses

The statistical analyses were performed using the Statistical Package for Social Sciences, version 16.0 (SPSS, Chicago, IL, USA). Results were expressed as means \pm SD, and a P value less than .05 was considered statistically significant. Normal distribution and homogeneity of the variances were tested. Parameters that did not fulfill normal distribution (IL-1 β , SBP, DBP, and monocytes count) were analyzed by nonparametric tests. Three individuals with outlier values for TNF- α concentration were excluded from TNF- α data analysis because the values were clearly extreme and do not belong to the general distribution. Data were analyzed using the Student t test (either paired or unpaired) and Mann-Whitney and Wilcoxon nonparametric tests, by comparing values during T2, T3, and T1. Correlation tests were performed using the Pearson parametric correlation test for normally distributed variables and Spearman nonparametric test for skewed variables.

3. Results

3.1. Changes in the blood pressure and anthropometric measurements

The mean values of SBP, DBP, body weight, BMI, body fat, WC, and HC obtained during T1, T2, and T3 are listed in Table 1.

Table 1 – Anthropometric and blood pressure characteristics of study subjects before, during, and after RIF (n = 50; mean age, 32.70 \pm 9.5 years)

Parameter	T1	T2	T3
BW (kg)	71.82 \pm 13.41	70.58 \pm 13.20***	71.92 \pm 13.50
BMI (kg/m ²)	26.30 \pm 5.01	25.85 \pm 4.91***	26.49 \pm 4.99
HC (cm)	102.61 \pm 9.66	101.90 \pm 9.98	101.02 \pm 9.67**
WC (cm)	83.62 \pm 11.17	82.69 \pm 10.34	82.68 \pm 10.37*
WC/HC ratio	0.81 \pm 0.07	0.81 \pm 0.07	0.81 \pm 0.06
BF (%)	24.12 \pm 12.60	20.38 \pm 11.32***	30.48 \pm 11.32***
SBP (mm Hg)	112.30 \pm 10.01	104.40 \pm 9.07***	111.82 \pm 9.60
DBP (mm Hg)	76.20 \pm 8.48	71.60 \pm 10.40**	74.50 \pm 14.40

Results are expressed as means \pm SD. BF, body fat; BW, body weight; T1, before Ramadan; T2, during Ramadan; T3, after Ramadan.

* P < .05, when statistically compared with T1 values.

** P < .01, when statistically compared with T1 values.

*** P < .001, when statistically compared with T1 values.

Compared with prefasting levels, Ramadan fasting resulted in a significant ($P < .001$) decrease in the body weight (70.58 ± 13.20 kg vs 71.82 ± 13.41 kg), BMI (25.85 ± 4.91 kg/m² vs 26.30 ± 5.01 kg/m²), and body fat percentage (20.38 ± 11.32 vs 24.12 ± 12.60). The latter was significantly increased (30.48 ± 11.32), however, after Ramadan (ie, cessation of fasting). In contrast, both WC and HC were slightly but not significantly decreased during Ramadan, and then they were significantly ($P < .05$ and $P < .01$, respectively) decreased after Ramadan, when compared with prefasting (T1). Comparing with prefasting levels (T1), both SBP and DBP were significantly ($P < .001$) decreased during Ramadan (T2) (104.40 ± 9.07 mm Hg vs 112.30 ± 10.01 mm Hg and 71.60 ± 10.40 mm Hg vs 76.20 ± 8.48 mm Hg, respectively) and then returned to the basal values measured 1 month after Ramadan (T3).

When considering the body weight changes during the study, a strong positive correlation ($P < .05$) was found between the body weight and the body fat percentage during the 3 study time points (Table 2).

3.2. Cytokines and immune cells

As revealed in Table 3 and compared with prefasting (T1) values, levels were significantly and substantially ($P < .001$) reduced during Ramadan for circulating IL-1 β (3.89 ± 4.84 pg/mL vs 17.84 ± 17.92 pg/mL), IL-6 (67.42 ± 51.25 pg/mL vs 155.85 ± 121.18 pg/mL), and TNF- α (52.22 ± 57.25 pg/mL vs 179.62 ± 129.56 pg/mL), with reduction values of about 78%, 57%, and 71%, respectively. However, IL-6 levels were maintained lower 1 month after Ramadan when compared with the basal levels (T1).

The total counts of leukocytes, granulocytes and, monocytes ($P < .001$) and lymphocytes ($P < .01$) were significantly decreased during Ramadan (T2) in comparison with before Ramadan (T1). Reduction values for WBC were less profound than those of the cytokines, and they were 14.1%, 14.4%, 11.2%, and 29.3%, respectively. Of a particular interest, the count of monocytes was significantly ($P < .01$) increased after Ramadan (T3) (Table 3).

Body weight, fat percentage, and caloric intake are among the most influencing factors that affect the levels of circulating cytokines. When considering the correlation between levels of circulating cytokines and those influencing factors during the 3 time points, we found that IL-1 β was negatively ($P < .05$) correlated with body fat percentage, whereas it was positively correlated ($P < .05$) with energy intake during the fasting stage (T2) (Table 4). However, for the 2 other cytokines, no significant correlations were found with any of the influencing factors during the fasting stage (T2), and IL-6 was positively correlated with energy intake during the

Table 3 – Circulating levels of proinflammatory cytokines and immune cells before, during, and after RIF (n = 50; mean age, 32.70 \pm 9.5 years)

Biomarker	T1	T2	T3
IL-6 (pg/mL) ^a	155.85 \pm 121.18	67.42 \pm 51.25***	109.6 \pm 75.6*
IL-1 β (pg/mL) ^a	17.84 \pm 17.92	3.89 \pm 4.84***	17.59 \pm 17.33
TNF- α (pg/mL) ^b	179.62 \pm 129.56	52.22 \pm 57.25***	147.62 \pm 120.07
Total leukocytes (10 ⁹ /L)	6.74 \pm 1.67	5.79 \pm 1.42***	6.51 \pm 1.91
Granulocytes (10 ⁹ /L)	4.02 \pm 1.37	3.44 \pm 1.20***	3.70 \pm 1.32
Lymphocytes (10 ⁹ /L)	2.32 \pm 0.71	2.06 \pm 0.52**	2.17 \pm 0.86
Monocytes (10 ⁹ /L)	0.41 \pm 0.15	0.29 \pm 0.10***	0.57 \pm 0.37**

Results are expressed as means \pm SD. T1, before Ramadan; T2, during Ramadan; T3, after Ramadan.

^a n = 40.

^b n = 48.

* $P < .05$, when statistically compared with T1 values.

** $P < .01$, when statistically compared with T1 values.

*** $P < .001$, when statistically compared with T1 values.

postfasting stage (T3), whereas TNF- α was positively correlated with energy intake during the prefasting stage (T1). It can be noticed, therefore, that energy intake was the most frequent factor that was positively correlated with the levels of the circulating cytokines (Table 4).

When considering the SBP and DBP, SBP was less intensely correlated with body weight and cytokine levels, whereas DBP was more profoundly correlated with these parameters. As shown in Tables 5 and 6, SBP was positively ($P < .05$) correlated with TNF- α only during the postfasting stage (T3), whereas DBP was positively ($P < .05$) correlated with IL-6, TNF- α , and body weight during the postfasting stage (T3) and with body weight only during the fasting stage (T2).

Interleukin-1 β is among the most important cytokines and first studied during Ramadan fasting in our study. Results of IL-1 β and its correlation with other cytokines and parameters (Table 7) revealed that it was positively correlated with IL-6 during the prefasting ($P < .001$) (T1) and postfasting ($P < .01$) (T3) stages, whereas it was positively correlated with TNF- α ($P < .05$) only during the postfasting stages (T3). Regarding its correlation with WBCs, IL-1 β was positively correlated with total leukocytes ($P < .01$) and granulocytes ($P < .05$) only during the prefasting stages. When considering the age as an influencing factor on IL-1 β , results revealed that age was positively correlated with IL-1 β during both prefasting (T1) ($P < .05$) and fasting (T2) stages.

3.3. Nutrient intakes

As shown in Table 8, results of nutrient analyses revealed that caloric energy, total protein, total carbohydrates, polyunsaturated fatty acids, trans-fat, cholesterol, α -Carotene, vitamin C, and lycopene did not differ significantly during Ramadan (T2) when compared with their prefasting levels (T1). However, water intake and folate ($P < .001$), total fat, saturated and monounsaturated fats ($P < .01$), and β -carotene ($P < .05$) intakes were significantly decreased during Ramadan (T2) when compared with prefasting (T1).

Table 2 – Association between BW and BF percentage before, during, and after RIF

	T1		T2		T3	
	Pearson r	P	Pearson r	P	Pearson r	P
Body fat (%)	0.380*	.007	0.438*	.001	0.445*	.001

Results are expressed as means \pm SD. BF, body fat; BW, body weight; T1, before Ramadan; T2, during Ramadan; T3, after Ramadan.

* $P < .05$ is considered statistically significant.

Table 4 – Associations between inflammatory cytokines, and BW, BF%, and EI

		T1			T2			T3		
		BW	BF (%)	EI	BW	BF%	EI	BW	BF%	EI
IL-1β	Pearson ρ	0.046	0.013	0.235	0.220	-0.37 *	0.344 *	0.092	0.086	0.271
	P	.783	.938	.203	.178	.018	.040	.579	.599	.100
IL-6	Pearson ρ	0.134	0.034	0.484	0.121	0.053	0.565	0.018	0.154	0.459 *
	P	.409	.835	.119	.455	.746	.098	.914	.343	.012
TNF-α	Pearson ρ	0.100	0.152	0.797 *	0.243	0.210	0.748	0.101	0.100	0.370
	P	.825	.318	.041	.107	.186	.051	.511	.514	.142

BF%, body fat percentage; BW, body weight; EI, energy intake; T1, before Ramadan; T2, during Ramadan; T3, after Ramadan.
 * P < .05 is considered statistically significant.

4. Discussion

It is now well recognized that long-lasting modifications in the circadian distribution of the eating and sleeping schedule during Ramadan fasting result in various changes in metabolism [26]. Significant weight loss and significant body fat reduction were observed during Ramadan and recovered to the baseline values 1 month after Ramadan. Our results were in accordance with previous findings showing a significant decrease in body weight [27,28] and a significant reduction in the body fat [29] during Ramadan. Because energy balance plays a regulatory role in body weight changes, some authors have associated Ramadan-related weight loss with the reduction in the total energy intake [22,28]. In our study, however, there was no significant difference in the total dietary energy intakes; even the difference between T1 and T2 was not a little amount of calories and equals the calorie of a small meal (555 kcal). Weight loss may also be explained by the negative fluid balance [30,31] because water intake significantly decreased among our subjects during Ramadan and then returned to baseline values 1 month after the end of Ramadan. The weight loss observed among our study subjects could be attributed to the reduced body fat because of the positive correlation between the 2 variables. Interestingly, although body fat was significantly increased after Ramadan, WC and HC were significantly decreased in the same time point. This may allow for the possibility of the pattern of body fat redistribution after Ramadan, which differs between males and females.

Given that TNF-α and IL-6 are well-known inhibitors of lipoprotein lipase (LPL) activity [32], decreased TNF-α and IL-6

levels may lead to increased LPL activity during Ramadan. This is in line with the increased LPL-catalyzed reaction products such as free fatty acids, during Ramadan [33]. Because glucose is less available during fasting, fat is considered a greater substrate for energy production, and accordingly, body fat and body weight are expected to be decreased [34]. This is in line with the insulin hyposecretion during Ramadan [22], favoring a predominant lipolytic state [33], which results in increased fat oxidation, observed during Ramadan [34].

Fasting reduces global cell proliferation rates [11,35], as evidenced in our study by the significant reduction in immune cells proliferation. Indeed, being in the reference ranges, total WBC, lymphocytes, monocytes, and granulocytes were significantly decreased during Ramadan and then returned to the basal values 1 month after Ramadan, except for monocytes, which significantly increased after Ramadan, which could be related to the rebound increase in postprandial activation as meal frequency increases after Ramadan [36]. These results were in accordance with the previous studies on healthy fasting volunteers [25,37,38] who showed a significant decrease in the total number of WBC during Ramadan.

In accordance with the antihypertensive effect of Ramadan fasting in previous studies [22,39,40], the present study also demonstrated decreases in both the SBP and the DBP. All participants were normotensive, which is expected given the fact that only 10 of them were older than 40 years. In fact, the decrease in blood pressure during Ramadan is more probably due to the reduction in body weight [39] or due to the inhibition of catecholamine production [41], which is known to be inhibited during hunger, causing a decrease in the

Table 5 – Associations between SBP and circulating cytokine levels and body weight before, during, and after RIF

Variable	T1		T2		T3	
	Spearman ρ	P	Spearman ρ	P	Spearman ρ	P
IL-1β	0.137	.400	0.017	.915	0.297	.096
IL-6	0.099	.544	0.088	.591	0.195	.199
TNF-α	0.036	.813	0.231	.126	0.351 *	.026
BW	0.047	.745	0.208	.147	0.164	.255

Results are expressed as means ± SD. BW, body weight; T1, before Ramadan; T2, during Ramadan; and T3, after Ramadan.
 * P < .05 is considered statistically significant.

Table 6 – Associations between DBP and levels of circulating cytokine and body weight before, during, and after RIF

Variable	T1		T2		T3	
	Spearman ρ	P	Spearman ρ	P	Spearman ρ	P
IL-1β	0.210	.192	0.040	.809	0.329 *	.038
IL-6	0.294	.066	0.007	.966	0.354 *	.025
TNF-α	0.028	.853	0.036	.825	0.168	.271
BW	0.134	.353	0.351 *	.012	0.305 *	.031

Results are expressed as means ± SD. BW, body weight; T1, before Ramadan; T2, during Ramadan; T3, after Ramadan.
 * P < .05 is considered statistically significant.

Table 7 – Associations of IL-1 β with other cytokines, anthropometric measures, WBCs, and age before, during, and after RIF

Variable	T1		T2		T3	
	Pearson ρ	P	Pearson ρ	P	Pearson ρ	P
TNF- α	0.293	.088	0.265	.124	0.429*	.010
IL-6	0.527***	.000	0.035	.228	0.422**	.007
BMI	0.056	.730	-0.308	.054	0.042	.796
BF%	0.013	.938	-0.372*	.018	0.086	.599
Total leukocytes	0.414**	.008	0.006	.972	0.198	.221
Lymphocytes	0.249	.122	0.148	.363	0.109	.502
Granulocytes	0.374*	.017	0.036	.825	0.177	.276
Monocytes ^a	0.137	.285	0.003	.986	0.120	.188
Age	0.378*	.016	0.423**	.007	0.047	.775

Results are expressed as means \pm SD. BF, body fat; T1, before Ramadan; T2, during Ramadan; and T3, after Ramadan.

^a Correlation test was performed using Spearman nonparametric test.

* $P < .05$.

** $P < .01$.

*** $P < .001$.

sympathetic tone and, as a consequence, falling of blood pressure, heart rate, and cardiac output. Our results agree with those that showed that IF has decreased blood pressure in rats [20,21]. Results of the current study are in agreement with those of Unalacak et al [24], who found that Ramadan fasting resulted in a significant reduction in SBP and DBP, as well WBC, TNF- α , and IL-2 and IL-8.

Interestingly, we found a positive association between SBP and TNF- α level after Ramadan. Similarly, a significant positive association was found between DBP and both IL-6 and IL-1 β after Ramadan. This is in consistent with the reported decrease in the plasma IL-6 level in association with the decreased SBP and DBP during Ramadan [42].

Unexpected negative correlation was found between IL-1 β levels and body fat percentage ($r = -0.372$, $P = .018$) during Ramadan. This is in contrast with previous findings linking adipose tissue to increased cytokine production, including IL-1 β , IL-6, and TNF- α [43].

More than 80% of cytokine production by adipose tissue originates from nonfat cells, of which macrophages are the major constituents [43]. Further, weight loss is associated with a reduction in macrophage infiltration of adipose tissue [44], and macrophages are mainly derived from monocytes. Accordingly, it could be postulated that the decreased cytokine levels may be related to the decrease in the number of monocytes during Ramadan, as reported in this study. An indirect support of this notion is the postprandial activation of monocytes expressing TNF- α and IL-1 β in response to meals [45] and the sustained increase in the monocyte secretion of TNF- α to more than 8 hours after each meal [36] and by the elevation of the number of monocytes expressing TNF- α within adipose tissue after a fat meal. Interestingly, the present study showed a significant positive correlation between IL-1 β and total WBC ($r = 0.414$, $P = .008$) and granulocytes ($r = 0.374$, $P = .017$), supporting the correlation between the cytokine level and the fasting-induced reduction in the number WBC.

The reduction in the cytokine level during Ramadan fasting could also be attributed to the decreased oxidative stress during Ramadan [46] and the decreased reactive oxygen species, which plays an important role in activating the transcriptional factor nuclear factor κ B [47], which is responsible for the expression of proinflammatory cytokines including IL-1 β , IL-6, and TNF- α [47], a matter that requires to be further investigated. Given that the antioxidant nutrients such as vitamin C, vitamin E, and carotenoids are involved in cytokines expression and secretion via activation of the nuclear factor κ B [47] and that our volunteer's diet during Ramadan (T2) was poor in these antioxidant nutrients including β -carotene and lycopene, the reductive effect of Ramadan fasting on the levels of proinflammatory cytokines

Table 8 – Dietary intake of different macronutrients and micronutrients by study subjects before, during, and after RIF (n = 50; mean age, 32.70 \pm 9.5 years)

Nutrient	T1	T2	T3
Energy (kcal)	2516.6 \pm 2144.98	1961.6 \pm 1165.69	2079.0 \pm 1446.04
Water (mL)	1585.4 \pm 884.10	1296.66 \pm 905.88***	1250.2 \pm 673.21
Total protein (g)	83.09 \pm 77.80	86.58 \pm 192.15	68.39 \pm 58.31
Total carbohydrates (g)	370.87 \pm 362.20	285.84 \pm 150.72	285.98 \pm 216.22
Total fat (g)	74.46 \pm 51.02	52.49 \pm 29.87**	71.78 \pm 68.64
Saturated fat (g)	17.50 \pm 14.61	12.34 \pm 8.33**	15.15 \pm 12.89
Monounsaturated fat (g)	23.12 \pm 20.55	14.03 \pm 11.50**	24.43 \pm 47.11
Polyunsaturated fat (g)	9.84 \pm 10.07	7.19 \pm 5.91	8.83 \pm 7.58
Trans-fat (g)	0.086 \pm 0.19	0.14 \pm 0.53	0.19 \pm 0.52
Cholesterol (mg)	205.39 \pm 236.48	193.73 \pm 373.51	130.98 \pm 172.18
α -Carotene (μ g)	79.53 \pm 358.77	87.83 \pm 347.93	255.71 \pm 865.43
β -Carotene (μ g)	616.09 \pm 1142.22	368.23 \pm 852.11*	892.65 \pm 2195.08
Vitamin C (mg)	59.95 \pm 52.42	74.72 \pm 85.95	87.67 \pm 150.20
Folate (μ g)	505.25 \pm 667.14	246.71 \pm 216.63***	358.31 \pm 445.81
Lycopene (μ g)	353.23 \pm 852.78	284.20 \pm 693.36	575.17 \pm 213.50

Results are expressed as means \pm SD. T1, before Ramadan; T2, during Ramadan; T3, after Ramadan.

* $P < 0.05$, when statistically compared with before T1 values.

** $P < .01$, when statistically compared with before T1 values.

*** $P < .001$, when statistically compared with before T1 values.

could not be ascribed to these elements but, rather, to the practice of IF and CR.

High-density lipoprotein (HDL) levels have been found to be increased during Ramadan fasting and maintained for at least 1 month after Ramadan [27,40], explaining, in our study, the sustained decrease in the IL-6 levels month after Ramadan. Although not statistically significant, TNF- α levels were also lowered 1 month after Ramadan. In addition, TNF- α interferes with lipid homeostasis and activates proatherogenic processes, including reduction of HDL cholesterol and increase of low-density lipoprotein cholesterol [48]. Although increased HDL levels during RIF were attributed to changes in dietary patterns [49], findings from our study raise the hypothesis that decreased IL-6 and TNF- α level might be the most probable explanation for the previously reported anti-atherogenic effect of Ramadan fasting [23]. The ability of RIF to reduce the levels of cytokines IL-6 and TNF- α upon fasting may drive a speculation that RIF could be beneficial for those with rheumatoid arthritis, a disease that had been reported to be characterized by an increased activity of the aforementioned cytokines [50], a matter that requires to be clinically investigated. Finally, because the 3 studied cytokines were found to be up-regulated in diabetes and reported to be significantly depressed in diabetic rats upon CR [51], it is worth to elaborate the impact of Ramadan fasting on diabetic people who practice Ramadan fasting, although those people are exempted from this worship and not allowed to fast.

This study maintains a few limitations. A potential limitation of this study was that, first, the protocol used in the present study may probably fail to give a better comparison between parameters in view of the prolonged investigation duration, which lasted from 10 AM to 4 PM for some, not all, of the subjects. Furthermore, it would be advisable to collect samples at 2 different time points each day (early morning and late afternoon) to elaborate the effect of fasting on 2 different periods of the day. Second, because both female and male subjects were included, the interruption of fasting during the menstrual period (5 ± 2 days) seems to affect the study results. However, data from female subjects showed a similar reduction in study parameters to that of male subjects. Third, the relatively small number of volunteers and, lastly, the lack of a control nonfasting group deprived the study from the ability to compare the effect of fasting to nonfasting, thus making it difficult to attribute the changes in cytokines and immune cells to the absolute effect of fasting.

Overall, our findings support the hypothesis that IF practiced during Ramadan by healthy subjects can effectively reduce inflammatory processes as evidenced by significantly reduced levels of leukocytes and circulating proinflammatory cytokines (IL-1 β , IL-6, and TNF- α). On the other hand, RIF effectively reduced body weight, body fat, and WC, and HC. From the results of the present study, it can be found out that although caloric intake was not significantly restricted upon fasting, cytokine levels were significantly reduced, indicating that practicing IF could be beneficial for health even it is not associated with significant CR. Furthermore, energy intake was the factor most frequently correlated with the changes of cytokines before, during, and after fasting, a matter that necessitates the significance of energy intake on the inflammatory process. This impact of Ramadan fasting on cytokines

provides biologically plausible mechanisms that may explain how Ramadan fasting may beneficially affect lipid and carbohydrate homeostasis as well as autoimmune diseases such as rheumatoid arthritis. The present study opens the door for future studies aiming to explore eventual mechanisms mediating this anti-inflammatory effect of RIF. It is suggested that further studies are to be done to elaborate the impact of RIF on patients with rheumatoid arthritis.

Acknowledgment

This research was financially supported by the Deanship of Graduate Studies and Scientific Research at Petra University (Grant 4/4/2009), Amman, Jordan. The authors are indebted to the Association of Agricultural Engineers of Jordan for supporting the researchers with the software of Food Processor SQL, for Mohammad Ehlayel for his thorough reviews, for Noor Hamed and Mona Al-Safadi for their technical assistance in nutritional assessment, and for Majed Hamdan for language reviewing.

Conflict of interest: None.

REFERENCES

- [1] Fontana L, Eagon JC, Trujillo ME, Scherer PE, Klein S. Visceral fat adipokine secretion is associated with systemic inflammation in obese humans. *Diabetes* 2007;56:1010–3.
- [2] You T, Yang R, Lyles MF, Gong D, Nicklas BJ. Abdominal adipose tissue cytokine gene expression: relationship to obesity and metabolic risk factors. *Am J Physiol Endocrinol Metab* 2005;288:741–7.
- [3] Petersen AM, Pedersen BK. The anti-inflammatory effect of exercise. *J Appl Physiol* 2005;98:1154–62.
- [4] Ugochukwu NH, Figgers CL. Dietary caloric restriction modifies inflammatory responses in livers of streptozocin-induced diabetic rats. *Nutr Res* 2006;26:221–6.
- [5] Lind L. Circulating markers of inflammation and atherosclerosis. *Atherosclerosis* 2003;169:203–14.
- [6] Spranger J, Kroke A, Möhlig M, Hoffmann K, Bergmann MM, Ristow M, et al. Inflammatory cytokines and the risk to develop type 2 diabetes: results of the prospective population-based European Prospective Investigation into Cancer and Nutrition (EPIC)-Potsdam Study. *Diabetes* 2006;52:812–7.
- [7] Sciarretta S, Ferrucci A, Ciavarella GM, De Paolis P, Venturelli V, Tocci G, et al. Markers of inflammation and fibrosis are related to cardiovascular damage in hypertensive patients with metabolic syndrome. *Am J Hypertens* 2007;20:784–91.
- [8] Federico A, D’Aiuto E, Borriello F, Barra G, Gravina AG, Romano M, et al. Fat: a matter of disturbance for the immune system. *World J Gastroenterol* 2010;16:4762–72.
- [9] Varady KA, Hellerstein MK. Alternate-day fasting and chronic disease prevention: a review of human and animal trials. *Am J Clin Nutr* 2007;86:7–13.
- [10] Mattson MP, Wan R. Beneficial effects of intermittent fasting and caloric restriction on the cardiovascular and cerebrovascular systems. *J Nutr Biochem* 2005;16:129–37.
- [11] Varady KA, Roohk DJ, McEvoy-Hein BK, Gaylinn BD, Thorne MO, Hellerstein MK. Modified alternate-day fasting regimens reduce cell proliferation rates to a similar extent as daily calorie restriction in mice. *FASEB J* 2008;22:2090–6.
- [12] Wan R, Ahmet I, Brown M, Cheng A, Kamimura N, Talan M, et al. Cardioprotective effect of intermittent fasting is

- associated with an elevation of adiponectin levels in rats. *J NutrBiochem* 2010;21:413–7.
- [13] Lu J, Lezi EE, Wang W, Frontera J, Zhu H, Wang WT, et al. Alternate day fasting impacts the brain insulin-signaling pathway of young adult male C57BL/6 mice. *J Neurochem* 2011;117:154–63.
- [14] Castello L, Froio T, Maina M, Cavallini G, Biasi F, Leonarduzzi G, et al. Alternate-day fasting protects the rat heart against age-induced inflammation and fibrosis by inhibiting oxidative damage and NF- κ B activation. *Free Radic Biol Med* 2010;48:47–54.
- [15] Brannon S, Gozansky W, Donahoo W, Melanson E, Cage C, Coussons-Read M. Obesity and inflammation: effects of short-term fasting on IL-6 and relationship to diurnal cortisol. *Bra BehImm* 2009;23:S28.
- [16] Aksungar FB, Topkaya AE, Akyildiz M. Interleukin-6, C-reactive protein and biochemical parameters during prolonged intermittent fasting. *Ann NutrMetab* 2007;51:188–95.
- [17] Johnson B, Summer W, Cutler RG, Martin B, Hyun DH, Dixit VD, et al. Alternate day calorie restriction improves clinical findings and reduces markers of oxidative stress and inflammation in overweight adults with moderate asthma. *Free Radic Biol Med* 2007;42:665–74.
- [18] Harvie MN, Pegington M, Mattson MP, Frystyk J, Dillon B, Evans G, et al. The effects of intermittent or continuous energy restriction on weight loss and metabolic disease risk markers: a randomized trial in young overweight women. *Int J Obes* 2011;35:714–27.
- [19] Arumugam TV, Phillips TM, Cheng A, Morrell CH, Mattson MP, Wan R. Age and energy intake interact to modify cell stress pathways and stroke outcome. *Ann Neurol* 2010;67:41–52.
- [20] Wan R, Camandola S, Mattson MP. Intermittent food deprivation improves cardiovascular and neuroendocrine responses to stress in rats. *J Nutr* 2003;133:1921–9.
- [21] Mager DE, Wan R, Brown M, Cheng A, Wareski P, Abernethy DR, et al. Caloric restriction and intermittent fasting alter spectral measures of heart rate and blood pressure variability in rats. *FASEB J* 2006;20:631–7.
- [22] Shariatpanahi ZV, Shariatpanahi MV, Shahbazi S, Hossaini A, Abadi A. Effect of Ramadan fasting on some indices of insulin resistance and components of the metabolic syndrome in healthy male adults. *Br J Nutr* 2008;100:147–51.
- [23] Aksungar FB, Eren A, Ure S, Teskin O, Ates G. Effects of intermittent fasting on serum lipid levels, coagulation status and plasma homocysteine levels. *Ann NutrMetab* 2005;49:77–82.
- [24] Unalacak M, Kara IH, Baltaci D, Erdem O, Bucaktepe PG. Effects of Ramadan fasting on biochemical and hematological parameters and cytokines in healthy and obese individuals. *Metab Syndr Relat Disord* 2011;9:157–61.
- [25] Deurenberg P, Deurenberg-Yap M, Guricci S. Asians are different from Caucasians and from each other in their body mass index/body fat per cent relationship. *Obes Rev* 2002;3:141–6.
- [26] Al Hourani HM, Atoum MF, Akel S, Hijawi S, Awawdeh S. Effects of Ramadan fasting on some hematological and biochemical parameters. *Jordan J BiolSci* 2009;2:103–8.
- [27] Adlouni A, Ghalim N, Saïle R, Hda N, Parra HJ, Benslimane A. Beneficial effect on serum apo AI, apo B and Lp AI levels of Ramadan fasting. *Clin Chim Acta* 1998;271:179–89.
- [28] Al-Numair K. Body weight and some biochemical changes associated with fasting in healthy Saudi men. *J Med Sci* 2006;6:112–6.
- [29] Al-Hourani HM, Atoum MF. Body composition, nutrient intake and physical activity patterns in young women during Ramadan. *Singapore Med J* 2007;48:906–10.
- [30] Gumaa KA, Mustafa KY, Mahmoud NA, Gader AM. The effects of fasting in Ramadan. 1. Serum uric acid and lipid concentrations. *Br J Nutr* 1978;40:573–81.
- [31] Sweileh N, Schnitzler A, Hunter GR, Davis B. Body composition and energy metabolism in resting and exercising Muslims during Ramadan fast. *J Sports Med Phys Fitness* 1992;32:156–63.
- [32] Coppack SW. Pro-inflammatory cytokines and adipose tissue. *ProcNutrSoc* 2001;60:349–56.
- [33] Chennaoui M, Desgorges F, Drogou C, Boudjema B, Tomaszewski A, Depiesse F, et al. Effects of Ramadan fasting on physical performance and metabolic, hormonal, and inflammatory parameters in middle-distance runners. *Appl Physiol Nutr Metab* 2009;34:587–94.
- [34] El Ati J, Beji C, Danguir J. Increased fat oxidation during Ramadan fasting in healthy women: an adaptative mechanism for body weight maintenance. *Am J Clin Nutr* 1995;62:302–7.
- [35] Varady KA, Roohk DJ, Hellerstein MK. Dose effects of modified alternate-day fasting regimens on in vivo cell proliferation and plasma insulin-like growth factor-1 in mice. *J ApplPhysiol* 2007;103:547–51.
- [36] Motton DD, Keim NL, Tenorio FA, Horn WF, Rutledge JC. Postprandial monocyte activation in response to meals with high and low glycemic loads in overweight women. *Am J ClinNutr* 2007;85:60–5.
- [37] Maughan RJ, Leiper JB, Bartagi Z, Zrifi R, Zerguini Y, Dvorak J. Effect of Ramadan fasting on some biochemical and hematological parameters in Tunisian youth soccer players undertaking their usual training and competition schedule. *J Sports Sci* 2008;3:39–46.
- [38] Sarraf-Zadegan N, Atashi M, Naderi GA, Baghai AM, Asgari S, Fatehifar MR, et al. The effect of fasting in Ramadan on the values and interrelations between biochemical, coagulation and hematological factors. *Ann Saudi Med* 2000;20:377–81.
- [39] Dewanti L, Watanabe C, Sulistiawati E, Ohtsuka R. Unexpected changes in blood pressure and hematological parameters among fasting and non fasting workers during Ramadan in Indonesia. *Eur J ClinNutr* 2006;60:877–81.
- [40] Mansi KMS. Study the effects of Ramadan fasting on serum glucose and lipid profile among healthy Jordanian students. *Am J App Sci* 2007;4:565–9.
- [41] Temizhan A, Dönderici O, Ouz D, Demirbas B. Is there any effect of Ramadan fasting on acute coronary heart disease events? *Int J Cardiol* 1999;70:149–53.
- [42] Fernandez-Real JM, Vayreda M, Richart C, Gutierrez C, Broch M, Vendrell J, et al. Circulating interleukin 6 levels, blood pressure, and insulin sensitivity in apparently healthy men and women. *J ClinEndocrinolMetab* 2001;86:1154–9.
- [43] Fain JN. Release of interleukins and other inflammatory cytokines by human adipose tissue is enhanced in obesity and primarily due to the nonfat cells. *Vitam Horm* 2006;74:443–77.
- [44] Bastard JP, Maachi M, Lagathu C, Kim MJ, Caron M, Vidal H, et al. Recent advances in the relationship between obesity, inflammation, and insulin resistance. *Eur Cytokine Netw* 2006;17:4–12.
- [45] Hyson DA, Paglieroni TG, Wun T, Rutledge JC. Postprandial lipemia is associated with platelet and monocyte activation and increased monocyte cytokine expression in normolipemic men. *Clin Appl Thromb Hemost* 2002;8:147–55.
- [46] Ibrahim WH, Habib HM, Jarrar AH, Al Baz SA. Effect of Ramadan fasting on markers of oxidative stress and serum biochemical markers of cellular damage in healthy subjects. *Ann Nutr Metab* 2008;53:175–81.
- [47] Portugal M, Barak V, Ginsburg I, Kohen R. Interplay among oxidants, antioxidants, and cytokines in skin disorders: present status and future considerations. *Biomed Pharmacother* 2007;61:412.
- [48] Fon Tracer KF, Kuzman D, Seliskar M, Pompon D, Rozman D. TNF- α interferes with lipid homeostasis and activates acute

- and proatherogenic processes. *Physiol Genomics* 2007;31: 216–27.
- [49] Chaouachi A, Chamari K, Roky R, Wong P, Mbazaa A, Bartagi Z, et al. Lipid profile of judo athletes during Ramadan. *Int J Sports Med* 2007;29:282–8.
- [50] Schuerwegh AJ, Dombrecht EJ, Stevens WJ, Van Offel JF, Bridts CH, De Clerck LS. Influence of pro-inflammatory (IL-1 α , IL-6, TNF- α , IFN- γ) and anti-inflammatory (IL-4) cytokines on chondrocyte function. *Osteoarthritis Cartilage* 2003;11:681–7.
- [51] Ugochukwu NH, Figgers CL. Caloric restriction inhibits up-regulation of inflammatory cytokines and TNF- α , and activates IL-10 and haptoglobin in the plasma of streptozotocin-induced diabetic rats. *J NutrBiochem* 2007;18: 120–6.